

# Neutrino Tridents at Near Detectors


Yuber F. Perez-Gonzalez

In collaboration with  
Peter Ballett - Matheus Hostert - Silvia Pascoli (Durham),  
Zahra Tabrizi and Renata Zukanovich Funchal (U-São Paulo)

Physics Opportunities in the Near  
DUNE Detector Hall  
Fermilab, Nov 2018



Based on arXiv:1807.10973




# Outline

- Introduction
- Trident Cross Section
- Events in LAr Detectors (DUNE ND)
- Conclusions

# Introduction

# Near Detector Program


Neutrino sector  
under scrutiny



- ◆ Flux
- ◆ CC and NC cross sections
- ◆ Backgrounds

# Near Detector Program

Neutrino sector  
under scrutiny



- ◆ Flux
- ◆ CC and NC cross sections
- ◆ Backgrounds

High beam luminosity +  
Large fiducial mass


Ideal places to investigate rare  
neutrino interactions

$$\sigma < 10^{-44} \text{ cm}^2$$

- ◆ Test SM predictions
- ◆ Search for BSM physics

# DUNE ND

## DUNE Beaml ine with Near Detector



Events per ton-year

$\nu_\mu$  CC Total  $1.64 \times 10^6$


$\nu_\mu$  NC Total  $5.17 \times 10^5$

$\nu_\mu - e$  135

Alan's talk

# DUNE ND

## DUNE Beaml ine with Near Detector



Events per ton-year

$\nu_\mu$  CC Total  $1.64 \times 10^6$

$\nu_\mu$  NC Total  $5.17 \times 10^5$

$\nu_\mu - e$  135


What about rare  
neutrino  
scatterings?

Alan's talk

# Trident Inelastic Scattering

Production of a charged lepton pair from the inelastic neutrino scattering in the Coulomb field of the nucleus


$$\nu_\alpha + \mathcal{H} \rightarrow \nu_{\alpha \text{ or } \kappa(\beta)} + \ell_\beta^- + \ell_\kappa^+ + \mathcal{H}$$



# Trident Inelastic Scattering

Production of a charged lepton pair from the inelastic neutrino scattering in the Coulomb field of the nucleus

$$\nu_\alpha + \mathcal{H} \rightarrow \nu_{\alpha \text{ or } \kappa(\beta)} + \ell_\beta^- + \ell_\kappa^+ + \mathcal{H}$$



# Trident Inelastic Scattering

Production of a charged lepton pair from the inelastic neutrino scattering in the Coulomb field of the nucleus

$$\nu_\alpha + \mathcal{H} \rightarrow \nu_{\alpha \text{ or } \kappa(\beta)} + \ell_\beta^- + \ell_\kappa^+ + \mathcal{H}$$

$$\nu_\mu \rightarrow \nu_\mu \mu^+ \mu^-$$

CHARM II

PLB 245 (1990) 271

$$\frac{\sigma_{\text{CHARM II}}}{\sigma_{\text{SM}}} = 1.58 \pm 0.57$$

CCFR


PRL 66 (1991) 3117

$$\frac{\sigma_{\text{CCFR}}}{\sigma_{\text{SM}}} = 0.82 \pm 0.28$$

NuTeV

$$\frac{\sigma_{\text{NuTeV}}}{\sigma_{\text{SM}}} = 0.67 \pm 0.27$$

Vancouver 1998, High energy physics, vol. 1



# Trident Inelastic Scattering

# Trident Inelastic Scattering

W. Altmannshoffer et. al., 2014

Z' constraints from trident

# Trident Inelastic Scattering

W. Altmannshoffer et. al., 2014

Z' constraints from trident

G. Magill and R. Plestid, 2016

Rates at DUNE ND

# Trident Inelastic Scattering

W. Altmannshoffer et. al., 2014

Z' constraints from trident

G. Magill and R. Plestid, 2016

Rates at DUNE ND

S.F. Ge et. al., 2016

Tridents produced by atmospheric neutrinos

# Trident Inelastic Scattering

W. Altmannshoffer et. al., 2014

Z' constraints from trident

G. Magill and R. Plestid, 2016

Rates at DUNE ND

S.F. Ge et. al., 2016

Tridents produced by atmospheric neutrinos

G. Magill and R. Plestid, 2017

Charged scalars

# Trident Inelastic Scattering

W. Altmannshoffer et. al., 2014

Z' constraints from trident

G. Magill and R. Plestid, 2016

Rates at DUNE ND

S.F. Ge et. al., 2016

Tridents produced by atmospheric neutrinos

G. Magill and R. Plestid, 2017

Charged scalars


Equivalent Photon Approximation (EPA)



Overestimates the cross section in some cases by more than 200%.

# Trident Cross Sections

$$\nu_\alpha(p_1) + \mathcal{H}(P) \rightarrow \nu_{\alpha \text{ or } \kappa(\beta)}(p_2) + \ell_\beta^-(p_4) + \ell_\kappa^+(p_3) + \mathcal{H}(P')$$



(Anti)Neutrino	SM Contributions
$\bar{\nu}_\mu \mathcal{H} \rightarrow \bar{\nu}_\mu \mu^- \mu^+ \mathcal{H}$	CC + NC
$\bar{\nu}_\mu \mathcal{H} \rightarrow \bar{\nu}_e e^\pm \mu^\mp \mathcal{H}$	CC
$\bar{\nu}_\mu \mathcal{H} \rightarrow \bar{\nu}_\mu e^- e^+ \mathcal{H}$	NC
$\bar{\nu}_e \mathcal{H} \rightarrow \bar{\nu}_e e^- e^+ \mathcal{H}$	CC + NC
$\bar{\nu}_e \mathcal{H} \rightarrow \bar{\nu}_\mu \mu^\pm e^\mp \mathcal{H}$	CC
$\bar{\nu}_e \mathcal{H} \rightarrow \bar{\nu}_e \mu^- \mu^+ \mathcal{H}$	NC

4-point limit

→ Observed!

$$V_{\alpha\beta\kappa}(A_{\alpha\beta\kappa}) \equiv g_V^\beta(g_A^\beta)\delta_{\beta\kappa} + \delta_{\alpha\beta}$$

Interference between  
CC and NC

# Trident Production

Differential  
cross sections

$$\frac{d^2\sigma_{\nu X}}{dQ^2 d\hat{s}} = \frac{1}{32\pi^2(s - M_{\mathcal{H}}^2)^2} \frac{H_X^{\mu\nu} L_{\mu\nu}}{Q^4}$$

# Trident Production

Differential  
cross sections

$$\frac{d^2\sigma_{\nu X}}{dQ^2 d\hat{s}} = \frac{1}{32\pi^2(s - M_{\mathcal{H}}^2)^2} \frac{H_X^{\mu\nu} L_{\mu\nu}}{Q^4}$$

Momentum Transfer 

$$s \equiv (p_1 + P)^2$$

Center-of-mass energy  
of the total system

$$\hat{s} \equiv (p_1 + q)^2$$

Center-of-mass energy  
of the neutrino-photon  
system

# Trident Production

Differential  
cross sections

$$\frac{d^2\sigma_{\nu X}}{dQ^2 d\hat{s}} = \frac{1}{32\pi^2(s - M_{\mathcal{H}}^2)^2} \frac{H_X^{\mu\nu} L_{\mu\nu}}{Q^4}$$

Hadronic Tensor

It depends on scattering regime

Momentum Transfer

$$s \equiv (p_1 + P)^2$$

Center-of-mass energy  
of the total system

$$\hat{s} \equiv (p_1 + q)^2$$

Center-of-mass energy  
of the neutrino-photon  
system

# Trident Production

Differential  
cross sections

It depends on scattering regime

Momentum Transfer

Hadronic Tensor

Leptonic Tensor

$$\frac{d^2\sigma_{\nu X}}{dQ^2 d\hat{s}} = \frac{1}{32\pi^2(s - M_{\mathcal{H}}^2)^2} \frac{Q^4}{Q^4}$$
$$H_X^{\mu\nu} L_{\mu\nu}$$

$$s \equiv (p_1 + P)^2$$

Center-of-mass energy  
of the total system

$$\hat{s} \equiv (p_1 + q)^2$$

Center-of-mass energy  
of the neutrino-photon  
system

# Trident Production

Differential cross sections

It depends on scattering regime

Momentum Transfer

Hadronic Tensor

Leptonic Tensor

$$\frac{d^2\sigma_{\nu X}}{dQ^2 d\hat{s}} = \frac{1}{32\pi^2(s - M_{\mathcal{H}}^2)^2} \frac{Q^4}{Q^4}$$

$H_X^{\mu\nu}$   $L_{\mu\nu}$

We can separate photon contributions

$$\frac{d^2\sigma_{\nu X}}{dQ^2 d\hat{s}} = \frac{1}{32\pi^2} \frac{1}{\hat{s} Q^2} \left[ h_X^T(Q^2, \hat{s}) \sigma_{\nu\gamma}^T(Q^2, \hat{s}) + h_X^L(Q^2, \hat{s}) \sigma_{\nu\gamma}^L(Q^2, \hat{s}) \right]$$

Transversal                      Longitudinal

$s \equiv (p_1 + P)^2$   
Center-of-mass energy  
of the total system

$\hat{s} \equiv (p_1 + q)^2$   
Center-of-mass energy  
of the neutrino-photon  
system

# Scattering Regimes

Coherent

# Scattering Regimes

Coherent


$$H_c^{\mu\nu} = 4Z^2e^2 \left| F(Q^2) \right|^2 \left( P^\mu - \frac{q^\mu}{2} \right) \left( P^\nu - \frac{q^\nu}{2} \right)$$

# Scattering Regimes

Coherent

$$H_c^{\mu\nu} = 4Z^2 e^2 \left| F(Q^2) \right|^2 \left( P^\mu - \frac{q^\mu}{2} \right) \left( P^\nu - \frac{q^\nu}{2} \right)$$

Cut on Q





# Scattering Regimes

CC      CC+NC      NC

Coherent

$$H_c^{\mu\nu} = 4Z^2 e^2 \left| F(Q^2) \right|^2 \left( P^\mu - \frac{q^\mu}{2} \right) \left( P^\nu - \frac{q^\nu}{2} \right)$$

Cut on Q





$$\sigma_0 = Z^2 \cdot 10^{-44} \text{ cm}^2$$

# Scattering Regimes

Diffractive

$$\frac{d^2\sigma_{\nu d}}{dQ^2 d\hat{s}} \rightarrow f(|\vec{q}|) \frac{d^2\sigma_{\nu d}}{dQ^2 d\hat{s}}$$

$$H_d^{\mu\nu}(P, P') = Z H_p^{\mu\nu}(P, P') + (A - Z) H_n^{\mu\nu}(P, P')$$





# Scattering Regimes

Diffractive

$$\frac{d^2\sigma_{\nu d}}{dQ^2 d\hat{s}} \rightarrow f(|\vec{q}|) \frac{d^2\sigma_{\nu d}}{dQ^2 d\hat{s}}$$

$$H_d^{\mu\nu}(P, P') = Z H_p^{\mu\nu}(P, P') + (A - Z) H_n^{\mu\nu}(P, P')$$



# Equivalent Photon Approximation (EPA)

Can we obtain an approximate cross section?

Fermi, 1924  
Weizsäcker, Williams, 1934

EPA

$$\sigma_t(P_i + C_s \rightarrow P_f + C_s) \approx \int dP(Q^2, \hat{s}) \sigma_\gamma(P_i + \gamma \rightarrow P_f; \hat{s}, Q^2 = 0)$$

# Equivalent Photon Approximation (EPA)


Can we obtain an approximate cross section?

Fermi, 1924  
Weizsäcker, Williams, 1934

EPA

Energy spectrum of  
the off-shell photons

$$\sigma_t(P_i + C_s \rightarrow P_f + C_s) \approx \int dP(Q^2, \hat{s}) \sigma_\gamma(P_i + \gamma \rightarrow P_f; \hat{s}, Q^2 = 0)$$



# Equivalent Photon Approximation (EPA)

Can we obtain an approximate cross section?

Fermi, 1924  
Weizsäcker, Williams, 1934

EPA

Energy spectrum of  
the off-shell photons

$$\sigma_t(P_i + C_s \rightarrow P_f + C_s) \approx \int dP(Q^2, \hat{s}) \sigma_\gamma(P_i + \gamma \rightarrow P_f; \hat{s}, Q^2 = 0)$$

Incoming particle  
with a real photon

# Equivalent Photon Approximation (EPA)

Can we obtain an approximate cross section?

Fermi, 1924  
Weizsäcker, Williams, 1934

EPA

Energy spectrum of  
the off-shell photons

$$\sigma_t(P_i + C_s \rightarrow P_f + C_s) \approx \int dP(Q^2, \hat{s}) \sigma_\gamma(P_i + \gamma \rightarrow P_f; \hat{s}, Q^2 = 0)$$

Assumptions

\* Longitudinal contribution can be neglected

$$\sigma_{\nu\gamma}^L(Q^2, \hat{s}) \approx 0$$

\* Transverse contribution can be taken as on-shell

$$\sigma_{\nu\gamma}^T(Q^2, \hat{s}) \approx \sigma_{\nu\gamma}^T(0, \hat{s})$$

## Assumptions

- \* Longitudinal contribution can be neglected

$$\sigma_{\nu\gamma}^L(Q^2, \hat{s}) \approx 0$$

- \* Transverse contribution can be taken as on-shell

$$\sigma_{\nu\gamma}^T(Q^2, \hat{s}) \approx \sigma_{\nu\gamma}^T(0, \hat{s})$$

## Trident case

Photon spectrum

$$\sigma_{\text{EPA}} = \int_{m_L^2}^{\hat{s}_{\max}} \int_{(\hat{s}/2E_\nu)^2}^{Q_{\max}^2} \sigma_{\nu\gamma}^T(0, \hat{s}) dP(Q^2, \hat{s})$$

$$dP(Q^2, \hat{s}) = \frac{Z^2 e^2}{4\pi^2} |F(Q^2)|^2 \frac{d\hat{s}}{\hat{s}} \frac{dQ^2}{Q^2}$$

## Assumptions

- \* Longitudinal contribution can be neglected

$$\sigma_{\nu\gamma}^L(Q^2, \hat{s}) \approx 0$$

- \* Transverse contribution can be taken as on-shell

$$\sigma_{\nu\gamma}^T(Q^2, \hat{s}) \approx \sigma_{\nu\gamma}^T(0, \hat{s})$$

## Trident case

$$\sigma_{\text{EPA}} = \int_{m_L^2}^{\hat{s}_{\max}} \int_{(\hat{s}/2E_\nu)^2}^{Q_{\max}^2} \sigma_{\nu\gamma}^T(0, \hat{s}) dP(Q^2, \hat{s})$$

Kinematical Limits

Photon spectrum

$$dP(Q^2, \hat{s}) = \frac{Z^2 e^2}{4\pi^2} |F(Q^2)|^2 \frac{d\hat{s}}{\hat{s}} \frac{dQ^2}{Q^2}$$

## Assumptions

- \* Longitudinal contribution can be neglected
- \* Transverse contribution can be taken as on-shell

$$\sigma_{\nu\gamma}^L(Q^2, \hat{s}) \approx 0$$

$$\sigma_{\nu\gamma}^T(Q^2, \hat{s}) \approx \sigma_{\nu\gamma}^T(0, \hat{s})$$

## Trident case

$$\sigma_{\text{EPA}} = \int_{m_L^2}^{\hat{s}_{\max}} \int_{(\hat{s}/2E_\nu)^2}^{Q_{\max}^2} \sigma_{\nu\gamma}^T(0, \hat{s}) dP(Q^2, \hat{s})$$

Photon spectrum

$$dP(Q^2, \hat{s}) = \frac{Z^2 e^2}{4\pi^2} |F(Q^2)|^2 \frac{d\hat{s}}{\hat{s}} \frac{dQ^2}{Q^2}$$

$$Q_{\text{cut}} = \frac{\Lambda_{\text{QCD}}}{A^{1/3}}$$

$$Q_{\text{cut}} < Q_{\max}$$

## Assumptions

- \* Longitudinal contribution can be neglected
- \* Transverse contribution can be taken as on-shell

$$\sigma_{\nu\gamma}^L(Q^2, \hat{s}) \approx 0$$

$$\sigma_{\nu\gamma}^T(Q^2, \hat{s}) \approx \sigma_{\nu\gamma}^T(0, \hat{s})$$

## Trident case

$$\sigma_{\text{EPA}} = \int_{m_L^2}^{\hat{s}_{\max}} \int_{(\hat{s}/2E_\nu)^2}^{Q_{\text{cut}}^2} \sigma_{\nu\gamma}^T(0, \hat{s}) dP(Q^2, \hat{s})$$

Photon spectrum

$$dP(Q^2, \hat{s}) = \frac{Z^2 e^2}{4\pi^2} |F(Q^2)|^2 \frac{d\hat{s}}{\hat{s}} \frac{dQ^2}{Q^2}$$

$$Q_{\text{cut}} = \frac{\Lambda_{\text{QCD}}}{A^{1/3}}$$

$$Q_{\text{cut}} < Q_{\text{max}}$$

## Assumptions

- \* Longitudinal contribution can be neglected
- \* Transverse contribution can be taken as on-shell

$$\sigma_{\nu\gamma}^L(Q^2, \hat{s}) \approx 0$$

$$\sigma_{\nu\gamma}^T(Q^2, \hat{s}) \approx \sigma_{\nu\gamma}^T(0, \hat{s})$$

## Trident case

$$\sigma_{\text{EPA}} = \int_{m_L^2}^{\hat{s}_{\max}} \int_{(\hat{s}/2E_\nu)^2}^{Q_{\text{cut}}^2} \sigma_{\nu\gamma}^T(0, \hat{s}) dP(Q^2, \hat{s})$$

Kinematical Limits

Photon spectrum

$$dP(Q^2, \hat{s}) = \frac{Z^2 e^2}{4\pi^2} |F(Q^2)|^2 \frac{d\hat{s}}{\hat{s}} \frac{dQ^2}{Q^2}$$

$$Q_{\text{cut}} = \frac{\Lambda_{\text{QCD}}}{A^{1/3}}$$

$$Q_{\text{cut}} < Q_{\text{max}}$$

Does it work?

# EPA

Does it work?

$$\sigma_t(P_i + C_s \rightarrow P_f + C_s) \approx \int dP(Q^2, \hat{s}) \sigma_\gamma(P_i + \gamma \rightarrow P_f; \hat{s}, Q^2 = 0)$$

# EPA

Does it work?

$$\sigma_t(P_i + C_s \rightarrow P_f + C_s) \approx \int dP(Q^2, \hat{s}) \sigma_\gamma(P_i + \gamma \rightarrow P_f; \hat{s}, Q^2 = 0)$$

QED

Fermi Limit of the SM

# EPA

Does it work?

$$\sigma_t(P_i + C_s \rightarrow P_f + C_s) \approx \int dP(Q^2, \hat{s}) \sigma_\gamma(P_i + \gamma \rightarrow P_f; \hat{s}, Q^2 = 0)$$

QED

Fermi Limit of the SM

$$\sigma_\gamma^{\text{QED}}(P_i + \gamma \rightarrow P_f; \hat{s}, 0) \propto \frac{1}{\hat{s}}$$

Decreases with  
increasing transferred  
four-momentum

On-shell  $\gg$  off-shell

# EPA

Does it work?

$$\sigma_t(P_i + C_s \rightarrow P_f + C_s) \approx \int dP(Q^2, \hat{s}) \sigma_\gamma(P_i + \gamma \rightarrow P_f; \hat{s}, Q^2 = 0)$$

QED

$$\sigma_\gamma^{\text{QED}}(P_i + \gamma \rightarrow P_f; \hat{s}, 0) \propto \frac{1}{\hat{s}}$$

Decreases with  
increasing transferred  
four-momentum

On-shell  $\gg$  off-shell

Fermi Limit of the SM

$$\sigma_\gamma^{\text{FL}}(P_i + \gamma \rightarrow P_f; \hat{s}, 0) \propto G_F^2 \hat{s}$$

Increases with  
increasing transferred  
four-momentum

On-shell  $\ll$  off-shell

# EPA

Does it work?

$$\sigma_t(P_i + C_s \rightarrow P_f + C_s) \approx \int dP(Q^2, \hat{s}) \sigma_\gamma(P_i + \gamma \rightarrow P_f; \hat{s}, Q^2 = 0)$$

QED

$$\sigma_\gamma^{\text{QED}}(P_i + \gamma \rightarrow P_f; \hat{s}, 0) \propto \frac{1}{\hat{s}}$$

Decreases with  
increasing transferred  
four-momentum

On-shell  $\gg$  off-shell

Fermi Limit of the SM

$$\sigma_\gamma^{\text{FL}}(P_i + \gamma \rightarrow P_f; \hat{s}, 0) \propto G_F^2 \hat{s}$$



Increases with  
increasing transferred  
four-momentum

On-shell  $\ll$  off-shell

EPA not valid for trident, right?

# EPA

EPA not valid for trident, right?




$$\frac{\sigma_{\nu\gamma}^T(Q^2, \hat{s})}{\sigma_{\nu\gamma}^T(0, \hat{s})} \quad \frac{\sigma^L(Q^2, \hat{s}) h_c^L(Q^2, \hat{s})}{\sigma^T(Q^2, \hat{s}) h_c^T(Q^2, \hat{s})}$$

Kinematically allowed region

$$E_\nu = 3 \text{ GeV}$$


# EPA

EPA not valid for trident, right?



Longitudinal contribution negligible

Transverse almost as on-shell



$$\frac{\sigma_{\nu\gamma}^T(Q^2, \hat{s})}{\sigma_{\nu\gamma}^T(0, \hat{s})} \quad \frac{\sigma^L(Q^2, \hat{s}) h_c^L(Q^2, \hat{s})}{\sigma^T(Q^2, \hat{s}) h_c^T(Q^2, \hat{s})}$$

Kinematically allowed region


$E_\nu = 3 \text{ GeV}$

# EPA

EPA not valid for trident, right?

Longitudinal contribution negligible

Transverse almost as on-shell




Physical cutoff on the momentum integration

$$\frac{\sigma_{\nu\gamma}^T(Q^2, \hat{s})}{\sigma_{\nu\gamma}^T(0, \hat{s})}$$

$$\frac{\sigma^L(Q^2, \hat{s}) h_c^L(Q^2, \hat{s})}{\sigma^T(Q^2, \hat{s}) h_c^T(Q^2, \hat{s})}$$

Kinematically allowed region



Form factor cuts the Q integration in a region which EPA works reasonably



$$E_\nu = 3 \text{ GeV}$$

# EPA

EPA not valid for trident, right?

# EPA

EPA not valid for trident, right?





# EPA

EPA not valid for trident, right?

Yes.

$$\mathcal{R} = \frac{\sigma_{\text{EPA}}(E_\nu)|_{Q_{\text{cut}}}}{\sigma_{\text{4PS}}(E_\nu)}$$



Compute the full 4PS cross sections

# Events in LAr Detectors

# Rates

$$N_{\text{X}}^{\psi} = \text{Norm} \times \int dE_{\nu} \sigma_{\nu\text{X}}(E_{\nu}) \frac{d\phi_{\nu}(E_{\nu})}{dE_{\nu}} \epsilon(E_{\nu})$$

# Rates

$$N_X^\psi = \text{Norm} \times \int dE_\nu \sigma_{\nu X}(E_\nu) \frac{d\phi_\nu(E_\nu)}{dE_\nu} \epsilon(E_\nu)$$

↓

$$\text{Exposure [POT]} \times \frac{\text{Fiducial Detector Mass} \times N_A}{m_T} [\text{target particles}]$$

Assuming 100% efficiency

# Rates

$$N_X^\psi = \text{Norm} \times \int dE_\nu \sigma_{\nu X}(E_\nu) \frac{d\phi_\nu(E_\nu)}{dE_\nu} \epsilon(E_\nu)$$

↓

$$\text{Exposure [POT]} \times \frac{\text{Fiducial Detector Mass} \times N_A}{m_T} [\text{target particles}]$$

Assuming 100% efficiency


Experiment	Baseline (m)	Total Exposure (POT)	Fiducial Mass (t)	$E_\nu$ (GeV)
SBND	110	$6.6 \times 10^{20}$	112	0 – 3
$\mu$ BooNE	470	$1.32 \times 10^{21}$	89	0 – 3
ICARUS	600	$6.6 \times 10^{20}$	476	0 – 3
DUNE	574	$12.81 (12.81) \times 10^{21}$	50	0 – 40
$\nu$ STORM	50	$1.83 \times (3 + 2 \times 2) \times 10^{21}$	100	0 – 6

↑  
 $\nu$  mode       $\bar{\nu}$  mode

$$\text{Exposure} = 1.83 \times (3 + 2 \times 2) \times 10^{21} \text{ POT}$$

# Rates

Higher cross  
sections at  
DUNE ND



# Rates


Exposure =  $1.83 \times (3 + 2 \times 2) \times 10^{21}$  POT

Channel	SBND	$\mu$ BooNE	ICARUS	DUNE ND	$\nu$ STORM ND
Not seen yet	Total $e^\pm\mu^\mp$	10 1	0.7 0.1	1 0.1	2993 (2307) 391 (299)
					191 23
Not seen yet	Total $e^+e^-$	6 0.2	0.4 0.0	0.7 0.02	1007 (800) 64 (49)
					114 6
Compare order of magnitudes	Total $\mu^+\mu^-$	0.4 0.3	0.0 0.0	0.0 0.0	286 (210) 143 (108)
					11 6

Coherent

Diffractive

Large contributions of diffractive events




# Kinematical Distributions



EPA gives also misleading distributions

Invariant charged lepton masses  $m_{\ell^+\ell^-}^2$

Small values



Peaked distributions




Tool for background suppression

Flux integrated distributions

All channels with same final states

# Kinematical Distributions

Separation angle  $\Delta\theta$



Flux integrated  
distributions

All channels with same  
final states

# Backgrounds?

Goal: Reach suppressions of order  $\mathcal{O}(10^{-6} - 10^{-5})$

- ♦ misID
 
$$\mu^+ \mu^- \longrightarrow \nu_\mu \text{CC}1\pi^\pm$$

$$e^+ e^- \longrightarrow \text{NC}\pi^0$$

$$e^\pm \mu^\mp \longrightarrow \text{CC}\pi^0$$

$$\text{CC}\gamma \qquad \nu_e \text{CC}\pi^\pm$$

misID	Rate
$\gamma$ as $e^\pm$	0.05
$\gamma$ as $e^+ e^-$	0.1 (w/ vertex)
$\pi^\pm$ as $\mu^\pm$	1 (no vertex + overlapping)


- ♦ No hadronic activity
- ♦  $m_{\mu^+ \mu^-}^2 < 0.2 \text{ GeV}^2, \Delta\theta < 20^\circ, \theta_\pm < 15^\circ$

A more careful analysis is needed

Channel	$\mathbf{N}_B^{\text{misID}}/\mathbf{N}_{\text{CC}}$	$\mathbf{N}_B^{\text{had}}/\mathbf{N}_{\text{CC}}$	$\mathbf{N}_B^{\text{kin}}/\mathbf{N}_{\text{CC}}$	$\epsilon_{\text{sig}}^{\text{coh}}$	$\epsilon_{\text{sig}}^{\text{dif}}$ 11
$e^\pm \mu^\mp$	$1.67 (1.62) \times 10^{-4}$	$2.68 (4.31) \times 10^{-5}$	$4.40 (3.17) \times 10^{-7}$	0.61 (0.61)	0.39 (0.39)
$e^+ e^-$	$2.83 (4.19) \times 10^{-4}$	$1.30 (2.41) \times 10^{-4}$	$6.54 (14.1) \times 10^{-6}$	0.48 (0.47)	0.21 (0.21)
$\mu^+ \mu^-$	$2.66 (2.73) \times 10^{-3}$	$10.4 (9.75) \times 10^{-4}$	$3.36 (3.10) \times 10^{-8}$	0.66 (0.67)	0.17 (0.16)



# Backgrounds?



# Conclusions

- A full 4PS computation is required to obtain correct estimates for the number of events.
- Let us stress that the EPA gives a reasonable result for the dimuon channel due to a serendipitous behavior of the Form Factor.
- EPA can artificially suppress the coherent scattering contribution and increase the diffractive one giving rise to an incorrect rate and distributions of observable quantities.
- We have estimated the background for each trident channel via a Monte Carlo simulation using GENIE, and identified the dominant contributions arising primarily from particle misidentification.
- Reduction of  $\sim 6$  orders of magnitude  $\times$  CC in the background is necessary to observe trident events at DUNE ND. A more careful analysis is needed.

# Thank you!





# Backup



# Coherent vs Diffractive

$$\sigma_{\nu\mathcal{N}} = \sigma_{\nu c} + \sigma_{\nu d}$$


$$\frac{\sigma_{\nu c}}{\sigma_{\nu\mathcal{N}}} \quad \frac{\sigma_{\nu d}}{\sigma_{\nu\mathcal{N}}}$$




# Rates




# Kinematical Distributions



# Kinematical Distributions



# Kinematical Distributions



# Events at other ND facilities

# Rates

$$N_X^\psi = \text{Norm} \times \int dE_\nu \sigma_{\nu X}(E_\nu) \frac{d\phi_\nu(E_\nu)}{dE_\nu} \epsilon(E_\nu)$$


Fraction

$$\sigma_{\nu X}^{\text{FAC}} = \sum_i f_i \sigma_{\nu X}^i$$


Cross section for the component

Experiment	Material	Baseline (m)	Exposure (POT)	Fiducial Mass (t)	$E_\nu$ (GeV)
INGRID	Fe	280	$3.9 \times 10^{21}$ [ $10^{22}$ ] T2K-I [T2K-II]	99.4	0 – 4
MINOS[+]	Fe and C	1040	$10.56(3.36)[9.69] \times 10^{20}$	28.6	0 – 20
NO $\nu$ A	$\text{C}_2\text{H}_3\text{Cl}$ and $\text{CH}_2$	1000	$8.85(6.9) [36(36)] \times 10^{20}$ [NO $\nu$ A-II]	231	0 – 20
MINER $\nu$ A	$\text{CH}, \text{H}_2\text{O}, \text{Fe}, \text{Pb}, \text{C}$	1035	$12(12) \times 10^{20}$	7.98	0 – 20

# Rates



# Rates



Channel	T2K-I	T2K-II	MINOS	MINOS+	NO $\nu$ A-I	NO $\nu$ A-II	MINER $\nu$ A
Total $e^\pm\mu^\mp$	563	1444	222 (56)	730	83 (72)	340 (374)	149 (102)
	96	246	46 (11)	151	25 (22)	102 (114)	56 (39)
Total $e^+e^-$	277	711	61 (15)	62	29 (22)	119 (114)	39 (27)
	24	62	9 (2)	8	4 (4)	16 (21)	10 (7)
Total $\mu^+\mu^-$	30	76	26 (6)	86	9 (9)	37 (47)	18 (13)
	21	54	15 (3)	49	8 (8)	34 (36)	18 (13)